

Coincidence Of An Insular Thyroid Carcinoma And Chronic Lymphocytic Leukemia; A Case Report

İnsular Tiroid Karsinomu ve Kronik Lenfositik Lösemi Birlikteliği: Olgu Sunumu

¹Doç.Dr. Mehmet YILMAZ

²Yrd.Doç.Dr. Güçhan ALANOĞLU

³Yrd.Doç.Dr. Nil GÜLER

⁴Yrd.Doç.Dr. S. Zeynep AKI

⁵Doç.Dr. Selda SEÇKİN

⁶Doç.Dr. Nuri Aydın KAMA

⁷Doç.Dr. Gülsüm ÖZET

¹Gaziantep University School of Medicine Department of Hematology

²Suleyman Demirel University School Of Medicine Department of Hematology

³19 Mayıs University School Of Medicine Department of Hematology

⁴Gazi University School Of Medicine Department of Hematology

⁵Ankara Numune Education and Research Hospital Department of Pathology

⁶Ankara Numune Education and Research Hospital Department of Surgery

⁷Ankara Numune Education and Research Hospital Department of Hematology

Gaziantep Tıp Dergisi 2010;16(1):15-18.

Özet

KLL olgun görünümlü eritrositlerin kan, karaciğer, kemik iliği, lenf nodları ve diğer dokularda birikimi ile karakterize hematolojik bir hastalıktır. Kronik lenfositik lösemide birlikte görülen hücrel ve humoral immün defektler nedeni ile ikincil maligniteler sık görülmekte ve enfeksiyona eğilim artmaktadır. Bu malignitelerden en sık görülenleri; büyük hücreli lenfoma (Richter Sendromu), Hodgkin hastalığı, akciğer ve cilt kanserleridir. Burada insular tiroid karsinomu ile kronik lenfositik lösemnin nadir birlikteliği sunulmuştur.

Anahtar kelimeler: KLL, İnsular karsinoma, İkincil malignansiler

Abstract

Chronic lymphocytic leukemia (CLL) is a hematological malignancy characterized with proliferation and accumulation of lymphocytes of mature appearance in blood, liver, bone marrow, lymph nodes and other tissues. Second malignancies occur with an increased frequency in CLL. The most frequent of these are large cell lymphoma, known as Richter's syndrome and Hodgkin's disease among hematological second malignancies, lung and skin cancers among solid tumors. The defect in humoral and cellular functions may be the underlying mechanism of increased tendency towards infections and second malignancies in CLL patients. Here we report a coexistence of insular thyroid carcinoma (ITC) and B-CLL in the same patient. The occurrence of both CLL and insular thyroid carcinoma described because of the rarity of the coexistence of these two disorders in the same patient.

Key Words: CLL, Thyroid, Insular carcinoma, Secondary malignancy

Introduction

Chronic lymphocytic leukemia (CLL) is a hematological malignancy characterized with proliferation and accumulation of lymphocytes of mature appearance in blood, liver, bone marrow, lymph nodes and other tissues. Secondary malignancies are frequent complications in patients with CLL with the standardized incidence ratio 1.59. (95% CI 1.50-1.69) (1).

Chemotherapeutic agents may also contribute to subsequent malignancies among CLL patients. Introduction of new treatment strategies including fludarabin was reported increased risk of lung cancer, acute myeloid leukemia and a more aggressive form of second malignancy (2). Alkylator-purine analog combination therapy may increase the risk of therapy-related myeloid malignancies (3). The secondary lymphoid malignancies complicating CLL includes prolymphocytic leukemia, Hodgkin disease, acute lymphoblastic leukemia, large B-cell non-Hodgkin lymphoma and multiple myeloma (4). Skin cancers, solid tumors and myeloproliferative disorders or myelodysplastic syndromes have also been documented (4-7).

The defect in humoral and cellular functions increases the tendency towards infections and secondary malignancies. Patients with chronic lymphocytic leukemia (CLL) may develop diffuse large B-cell lymphoma (DLBL), also known as Richter's syndrome. Richter's syndrome may develop in both mutated and unmutated immunoglobulin heavy-chain variable region genes of CLLs, but clonal transformation of CLL to DLBL occur only in the unmutated subgroup of CLL (8). Here we report a coexistence of an insular thyroid carcinoma and CLL.

Doç.Dr. Mehmet YILMAZ, Gaziantep Üniversitesi Tıp Fakültesi, İç Hastalıkları AD, Hematoloji BD.

Adres: Gaziantep Üniversitesi Tıp Fakültesi, Şehitkamil/GAZİANTEP

Tel: 0342 360 60 60 **Dahili:** 76128/76140 **E-mail:** mnyilmaz@gantep.edu.tr

Geliş Tarihi: 26.10.2009 **Kabul Tarihi:** 23.12.2009



Case Report

A 50-year-old man referred to Ankara Numune Education and Research Hospital hematology department because of a swelling on the anterior side of his neck enlarging during the last two years. On physical examination, 15x15 cm in diameter clover sized mass which consists of combined masses on the localization of thyroid gland; multiple lymphadenopathies in submandibular, inguinal and axillary region varying in size, a mass of 1x2 cm in diameter on the sixth and seventh costosternal junction and splenomegaly were detected. Laboratory values were WBC 63.200/ μ L, lymphocyte 58.000/ μ L, Hb 11.1 g/dL, and b2 micro globulin 13 mgr/L (N: 1-1, 7).

Thyroid function tests and thyroid antibodies were normal as were the platelet counts, ESR and LDH. There were mature lymphocytes and basket cells on blood smear. Bone marrow aspiration disclosed lymphocyte cell lines of 80-85 percent (Figure 3, 4). Flow cytometric analysis showed CD20 (95%), CD23 (72%), and CD19/CD5 together (40%) on lymphocytes. IgM-kappa surface immunoglobulins on B-cell lines were also detected. Findings of bone marrow examination and flow cytometry analysis were consistent with stage II CLL according to RAI Clinical Staging System. A chest tomography disclosed multiple metastatic lesions in the lungs, mediastinal lymphadenopathies and a mass on the 6th, 7th, and 8th costae destructing the bone structure. The histopathological examination of the biopsies from the mass localized on the thyroid gland showed poorly differentiated insular carcinoma (Figure 1).

Histopathological examinations of the cervical, submandibular lymph node sections revealed the infiltrations of the tumor cells on lymph node capsule and extracapsullary field. Immunohistochemical studies showed the positive stained cells with LCA, CD-20 and lambda chain findings consistent with CLL (Figure 2). Besides, metastases of insular thyroid carcinoma on submandibular and cervical lymph nodes were seen. Thus, he underwent an operation of bilaterally thyroidectomy and debulking.

Discussion

Insular thyroid carcinoma was an interval form between well differentiated (papillary and follicular) and poorly differentiated carcinoma (anaplastic). It stems from the epithelium of thyroid follicles (9). Recently, it has been classified as a distinct entity. Thyroid insular carcinoma can develop on the ground of a goiter over in a long period. William IK et al (10) described a case of having huge goiter 2 year prior to the diagnosis of an insular thyroid carcinoma which extends through substernally. This may be inferred by proliferation, mutation, and transformation of the tumor physiopathology.




Figure 1. Tumor cells composing solid islets adjacent to thyroid tissue (X40 H&E).




Figure 2. Lymphocytes of uniform shaped having hyperchromatic small cytoplasm in the lymph nodes adjacent to thyroid tissue (x400 H&E).




Figure 3. Lymphocytes diffusely infiltrated in bone marrow (X10, Giemsa)




Figure 4. Lymphocytes of uniform shaped, mature appearance and hyperchromatic nucleus scanty cytoplasm in bone marrow aspiration (X100, Giemsa)

Moreover, in a 22-patient study, it is reported that 27 percent of these patients having insular thyroid carcinoma had a goiter history of 10 years or more. (11). In our case he had 25-year goiter history. There was no exposure to radiation or I-131 intake and metastasis to lungs and bones were detected. Biopsy taken from the right sixth costae revealed the metastasis of an insular thyroid carcinoma. Konturek et al. alleged to be similarities between the chromosomal changes in the thyroid carcinomas and those of the changes in breast carcinomas and CLL (12). Evarus et al. have determined dysfunction in B and T lymphocytes in early stage CLL, regulation disorder in concurrence hypothalamic-pituitary-adrenal axis, disturbance in sex hormones and changes in thyroid hormones. According to this they claimed that the disturbance in immunohormonal interactions in early-stage CLL were responsible for the pathogenesis of in this lymphoproliferative malignancy (13). Immune disturbances were accompanied by hormonal imbalance, depending on disease status: lower ACTH, cortisol and triiodothyronine levels were established to occur in active CLL compared to indolent disease (14). But, Paydas et al (15) showed an association between 7 patients previously having been diagnosed as chronic lymphoproliferative disease and thyrotoxicosis.

CLL is a classical form of human malignancy that arises from the defects of apoptosis. It occurs by the aberrations of the genes that regulate this apoptosis (16). Second malignancies in CLL occur with an increased frequency mainly result of the immune defects associated with this disease. An antigenic stimulation through the B-cell receptor is involved in the selection and possibly also the expansion of the malignant clone. Though all evidences suggest specific Ag recognition and possibly stimulation at different time-points, the nature of the Ag(s) is still unknown. It appears likely that CLL cells derive from a pool of auto/polyreactive CD5(+) B cells. (17,18).

Although among the patients having been followed with a diagnosis of CLL the risk for the development of secondary malignancies is high. No case has been reported to be diagnosed concurrently as CLL and Insular thyroid carcinoma in the literature. Likely Immunologic defects to CLL may be involved in this presentation of disease.

In conclusion, this occurrence of both CLL and insular thyroid carcinoma described because of the rarity of the coexistence of these two diseases in the same patient.

References

- Schöllkopf C, Rosendahl D, Rostgaard K. Risk of second cancer after chronic lymphocytic leukemia. *Int J Cancer*. 2007;121:151-156.
- Meloni G, Proia A, Guerrisi V, Cordone I. Acute myeloid leukemia and lung cancer occurring in a chronic lymphocytic leukemia patient treated with fludarabine and autologous stem cell transplantation. *Ann Oncol*. 2000;11:1493-1495.
- Morrisan VA, Rai KR, Peterson BL. Therapy-related myeloid leukemias are observed in patients with chronic lymphocytic leukemia after treatment with fludarabine and chlorambucil: results of an intergroup study, cancer and leukemia group B 9011. *J Clin Oncol*. 2002;20:3878-3884.
- Copur MS, Ledakis P, Novinski D. An unusual case of composite lymphoma involving chronic lymphocytic leukemia, follicular lymphoma and Hodgkin disease. *Leuk and Lymphoma*. 2004;45:1071-1076.
- Hisada M, Biggar RJ, Greene MH. Solid Tumor after chronic lymphocytic leukemia. *Blood*. 2001;96:1979-81.
- Gargallo P, Cacchione R, Chena C. Chronic lymphocytic leukemia developing in a patient with chronic myeloid leukemia: evidence of distinct lineage-associated genomic events. *Cancer Genet Cytogenet*. 2005;161:74-77.
- Varkonyi J, Zalatnai A, Timar J. Secondary cutaneous infiltration in B cell chronic lymphocytic leukemia. *Acta Haematol*. 2000;103:116-121.
- Timar B, Fülöp Z, Csernus B. Relationship between the mutational status of VH genes and pathogenesis of diffuse large B-cell lymphoma in Richter's Syndrome. *Leukemia*. 2004;18:326-330.
- Sironi M, Collini P, Cantaboni A. Fine needle aspiration cytology of insular thyroid carcinoma, A report of four cases. *Acta Cytologica*. 1992;36:435-39.
- Kuhel WI, Kutler DI, Santos-Buch CA. Poorly differentiated insular thyroid carcinoma, a case report with identification of intact insulae with fine needle aspiration biopsy. *Acta Cytologica*. 1998;42:991-997.
- Lam KY, Lo CY, Chan KW. Insular and anaplastic carcinoma of the thyroid: a 45-year comparative study at a single institution and review of the significance of p53 and p21. *Ann Surg*. 2000;231:329-38.
- Konturek A, Barczynski M, Kostyk E. Chromosome aberrations in patients with papillary thyroid cancer and other neoplasms. *Przegl Lek*. 1998;55:290-293.
- Everaus H, Lehtmaa J, Luik E. Immune and hormonal changes in early-stage chronic lymphocytic leukemia patients. *Ann Hematol*. 1992;65:219-23.
- Evaraus H, Luik E, Lehtmaa J. Active and indolent chronic lymphocytic leukaemia--immune and hormonal peculiarities. *Cancer Immunol Immunother*. 1997;45:109-114.

15. Paydaş S, Sahin B, Hazar B. The association of thyrotoxicosis and chronic lymphoproliferative disorders. *Leuk Lymphoma*. 2000;38:131-135.
16. Kips Tj. Signal transduction pathways and mechanisms of apoptosis in CLL-B lymphocytes: their role in CLL pathogenesis. *Hematol Cell Ther*. 1997;39:17-27.
17. Conley CI, Misiti J, Laster AJ. Genetic factors predisposing to chronic lymphocytic leukemia and to auto-immun disease. *Medicine*. 1980;59:323-334.
18. Ghia P, Scielzo C, Frenquelli M, Muzie, M. From normal to clonal B cells: Chronic lymphocytic leukemia (CLL) at the crossroad between neoplasia and autoimmunity. *Autoimmun Rev*. 2007;7:127-131.